

Spectral Properties and Linear Stability of Self-Similar Wave Maps

Roland Donniger* and Peter C. Aichelburg†

Faculty of Physics, Gravitational Physics
University of Vienna
Boltzmannngasse 5, A-1090 Wien

August 1, 2009

Abstract

We study co-rotational wave maps from $(3 + 1)$ -Minkowski space to the three-sphere S^3 . It is known that there exists a countable family $\{f_n\}$ of self-similar solutions. We investigate their stability under linear perturbations by operator theoretic methods. To this end we study the spectra of the perturbation operators, prove well-posedness of the corresponding linear Cauchy problem and deduce a growth estimate for solutions. Finally, we study perturbations of the limiting solution which is obtained from f_n by letting $n \rightarrow \infty$.

1 Introduction and Definition of the Model

Many nonlinear evolution equations show a behaviour known as "blow up": Although the initial data are regular, the solution becomes singular in finite time. Important questions in this respect are: Under what circumstances does the blow up occur? What is the mechanism that leads to singularity formation? What is the blow up rate and the shape of the profile before the singularity forms? Unfortunately, many interesting examples from physics (such as Einstein's equations) are much too complicated to be accessible to presently available analytical techniques. Thus, one has to rely on numerical studies and simplified "toy models" which share some features with the original equation but are easier to handle.

The wave maps system provides an example of such a toy model which has been studied in recent times. Consider $(n + 1)$ -dimensional Minkowski space (\mathbb{R}^{n+1}, η) with metric $(\eta_{\mu\nu}) = \text{diag}(-1, 1, \dots, 1)$, an m -dimensional Riemannian manifold (M, g) with metric g and a mapping $\Phi : \mathbb{R}^{n+1} \rightarrow M$. Choosing local coordinates on M we denote the components of Φ and the metric g by Φ^A and g_{AB} , respectively, where capital latin indices run from 1 to m . The action functional S for wave maps is defined by

$$S(\Phi) = \int_{\mathbb{R}^{n+1}} \eta^{\mu\nu} (\partial_\mu \Phi^A) (\partial_\nu \Phi^B) g_{AB} \circ \Phi$$

where integration is understood with respect to the ordinary Lebesgue measure on \mathbb{R}^{n+1} and Einstein's summation convention is in force (greek indices run from 0 to n). As usual we write $\partial_\mu := \frac{\partial}{\partial x^\mu}$. The Euler-Lagrange equations associated to this action are given by

$$\square \Phi^A + \eta^{\mu\nu} \Gamma^A_{BC}(\Phi) (\partial_\mu \Phi^B) (\partial_\nu \Phi^C) = 0 \quad (1)$$

where $\Gamma^A_{BC} := \frac{1}{2} g^{AD} (\partial_B g_{CD} + \partial_C g_{BD} - \partial_D g_{BC})$ are the Christoffel symbols associated to the metric g on M and $\square := \eta^{\mu\nu} \partial_\mu \partial_\nu$ denotes the wave operator. Eq. (1) is known as the *wave maps equation* and it is a system of semilinear wave equations.

We are interested in the Cauchy problem for eq. (1), i.e. we prescribe initial data $\Phi^A|_{t=0} = \Phi_0^A$ and $\partial_t \Phi^A|_{t=0} = \Phi_1^A$ where $\partial_t := \partial_0$ and study the future development. There exists a well-established local

*roland.donniger@univie.ac.at

†aichelb8@univie.ac.at

well-posedness theory for the wave maps system which states that for any initial data $(\Phi_0^A, \Phi_1^A) \in H^s \times H^{s-1}$ ($s > \frac{n}{2}$) there exists a $T > 0$ and functions $\Phi^A \in C([0, T], H^s)$ with $\partial_t \Phi^A \in C([0, T], H^{s-1})$ that solve eq. (1) in the sense of distributions and satisfy $(\Phi^A, \partial_t \Phi^A)|_{t=0} = (\Phi_0^A, \Phi_1^A)$ where $H^s := H^s(\mathbb{R}^n)$ denotes the usual fractional Sobolev space. Moreover, the time T can be chosen to depend continuously on the initial data and the mapping $(\Phi_0^A, \Phi_1^A) \mapsto (\Phi^A, \partial_t \Phi^A) : H^s \times H^{s-1} \rightarrow C([0, T], H^s) \times C([0, T], H^{s-1})$ is continuous (see [6] and references therein for details). There are also many recent results addressing global issues, see e.g. [16] for a review. Of most interest for our purposes is Tao's work [15] which deals with spherical targets, i.e. M is an $(m-1)$ -sphere for $m, n \geq 2$. Under this assumption a global smooth solution of the wave maps system eq. (1) exists whenever the initial data are smooth and small in the homogeneous Sobolev space $\dot{H}^{n/2} \times \dot{H}^{n/2-1}$.

The present work deals with co-rotational wave maps from $(3+1)$ -Minkowski space to the three-sphere S^3 . Choosing the usual spherical coordinates (t, r, θ, φ) on Minkowski space and standard coordinates (ψ, Θ, ϕ) on S^3 with metric $g = d\psi^2 + \sin^2 \psi (d\Theta^2 + \sin^2 \Theta d\phi^2)$, we restrict ourselves to mappings of the form $(t, r, \theta, \varphi) \mapsto (\psi(t, r), \theta, \varphi)$ which are called *co-rotational*. Under these assumptions the system eq. (1) reduces to the single semilinear wave equation

$$\psi_{tt} - \psi_{rr} - \frac{2}{r} \psi_r + \frac{\sin(2\psi)}{r^2} = 0. \quad (2)$$

We remark that Tao's result applies to this problem but there are earlier papers by Sideris [13] and Shatah, Tahvildar-Zadeh [12] which address global questions for this particular model.

2 Self-Similar Solutions and Singularity Formation

Note that eq. (2) is invariant under dilations, i.e. if ψ is a solution, so is ψ_λ defined by $\psi_\lambda(t, r) := \psi(t/\lambda, r/\lambda)$ for a $\lambda > 0$. There exists a conserved energy associated to eq. (2) given by

$$E_\psi(t) = \int_0^\infty (r^2 \psi_t^2(t, r) + r^2 \psi_r^2(t, r) + 2 \sin^2(\psi(t, r))) dr$$

that scales as $E_{\psi_\lambda} = \lambda E_\psi$ which means that eq. (2) is *energy supercritical*. It is widely believed that for energy supercritical equations blow up is possible since it is energetically favourable for solutions to shrink which might result in the formation of a singularity. In order to obtain an explicit example of a blow up solution it is reasonable to look for self-similar solutions $\psi(t, r) = f(\frac{r}{T-t})$ where $T > 0$ is a parameter (the *blow up time*). Plugging this ansatz into eq. (2) yields

$$f'' + \frac{2}{r} f' - \frac{\sin(2f)}{\rho^2(1-\rho^2)} = 0 \quad (3)$$

where $' := \frac{d}{d\rho}$ and $\rho := \frac{r}{T-t}$. We require the regularity conditions $f(0) = 0$ and $f(1) = \frac{\pi}{2}$. Shatah [11] has shown existence of a smooth solution f_0 of eq. (3) which has later been found in closed form [17] and it is given by $f_0(\rho) = 2 \arctan(\rho)$. Bizoń [2] has extended this result by proving existence of a countable family $\{f_n : n = 0, 1, \dots\}$ of analytic solutions of eq. (3) satisfying $f_n(0) = 0$ and $f_n(1) = \frac{\pi}{2}$ where n counts the number of intersections of f_n with the line $\frac{\pi}{2}$ on the interval $[0, 1)$. Furthermore, the solutions f_n converge to the constant $f_\infty \equiv \frac{\pi}{2}$ for $n \rightarrow \infty$ pointwise on $(0, 1]$. These self-similar wave maps provide explicit examples of solutions with smooth finite energy initial data (use appropriate cut-off functions and finite speed of propagation) that become singular when $t \rightarrow T^-$.

An obviously important question is whether blow up occurs for generic initial data or the self-similar examples are "exceptional cases". This issue has been addressed by Bizoń et. al. [4] using numerical techniques. Their results strongly suggest that the self-similar solution f_0 provides a universal blow up profile. More precise, they conjecture that there exists a large set of initial data that lead to blow up and the singularity formation takes place via f_0 , i.e. the time evolution approaches f_0 near the center $r = 0$ as $t \rightarrow T^-$.

We remark that this self-similar blow up behaviour is very different from singularity formation in the energy critical case of co-rotational wave maps from $(2+1)$ -Minkowski space to the two-sphere. For this model it has been known for quite some time (see [14]) that the blow up rate cannot be self-similar. However, rigorous existence of blow up solutions has only recently been shown by Rodnianski and Sterbenz [10] as well as Krieger, Schlag, Tataru [7].

3 The Linear Stability Problem

As mentioned, the self-similar solution f_0 seems to play an important role in dynamical time evolution. Thus, it has to be stable in a certain sense since otherwise it could not be found by numerical time evolutions starting from generic initial data. In order to study linear stability of the wave map f_n we introduce adapted coordinates (σ, ρ) given by $\sigma = -\log \sqrt{(T-t)^2 - r^2}$ and $\rho = \frac{r}{T-t}$ (cf. [2]). The introduction of similarity coordinates in the study of self-similar blow up is very common (see e.g. [8]). We refer to (σ, ρ) as *hyperbolic coordinates* since the lines $\sigma = \text{const}$ are hyperbolae in a spacetime diagram. Note that the coordinate system (σ, ρ) is only defined for $r < T - t$ and thus, it covers the interior of the backward lightcone of the blow up point $(t, r) = (T, 0)$. Furthermore, (σ, ρ) is an orthogonal coordinate system, i.e. the vectorfields ∂_σ and ∂_ρ are orthogonal with respect to the Minkowski metric. Moreover, the blow up time $t = T$ is shifted to infinity, i.e. $t \rightarrow T-$ corresponds to $\sigma \rightarrow \infty$. The valid range is $\sigma \in \mathbb{R}$ and $\rho \in [0, 1)$. Eq. (2) transforms into

$$\psi_{\sigma\sigma} - 2\psi_\sigma - (1 - \rho^2)^2 \psi_{\rho\rho} - \frac{2(1 - \rho^2)^2}{\rho} \psi_\rho + \frac{(1 - \rho^2) \sin(2\psi)}{\rho^2} = 0. \quad (4)$$

Plugging the ansatz $\psi(\sigma, \rho) = f_n(\rho) + \phi(\sigma, \rho)$ into eq. (4) and linearizing around f_n we arrive at the evolution equation

$$\phi_{\sigma\sigma} - 2\phi_\sigma - (1 - \rho^2)^2 \phi_{\rho\rho} - \frac{2(1 - \rho^2)^2}{\rho} \phi_\rho + \frac{2(1 - \rho^2) \cos(2f_n)}{\rho^2} \phi = 0 \quad (5)$$

for the perturbation ϕ which governs the linearized flow around the wave map f_n . The wave map f_n is considered to be linearly stable if there do not exist solutions of eq. (5) that grow (with respect to a suitable norm) as σ increases. We will make this more precise below. However, a direct computation shows that ϕ_n^G defined by $\phi_n^G(\sigma, \rho) := e^\sigma \rho \sqrt{1 - \rho^2} f_n'(\rho)$ solves eq. (5). This instability is related to the time translation symmetry of the original problem. The reason for this is the fact that f_n is not a single solution but a one-parameter family of solutions depending on the blow up time T . We will refer to this instability as the *gauge instability*. Thus, without further considerations the best we can expect is to prove a result that excludes the existence of solutions of eq. (5) with $f_n = f_0$ that grow *faster* than this gauge instability. In what follows we will address this issue by using an operator theoretic approach.

4 Operator Formulation

In order to simplify the problem further we define a new unknown $\tilde{\phi}$ by $\tilde{\phi}(\sigma, \rho) := e^{-\sigma} \phi(\sigma, \rho)$ to remove the first-order term $-2\phi_\sigma$. Eq. (5) transforms into

$$\tilde{\phi}_{\sigma\sigma} - (1 - \rho^2)^2 \tilde{\phi}_{\rho\rho} - \frac{2(1 - \rho^2)^2}{\rho} \tilde{\phi}_\rho + \frac{2(1 - \rho^2) \cos(2f_n) - \rho^2}{\rho^2} \tilde{\phi} = 0. \quad (6)$$

4.1 The Operator A_n

We define the formal Sturm-Liouville differential expression a_n on the interval $(0, 1)$ by

$$a_n u := \frac{1}{w} (-(pu)') + q_n u$$

where $w(\rho) := \frac{\rho^2}{(1-\rho^2)^2}$, $p(\rho) := \rho^2$ and $q_n(\rho) := \frac{2(1-\rho^2) \cos(2f_n(\rho)) - \rho^2}{(1-\rho^2)^2}$. Thus, a_n represents the spatial differential operator in eq. (6). We define the weighted Hilbert space $H := L_w^2(0, 1)$ where $L_w^2(0, 1)$ consists of (equivalence classes of) functions which are square integrable with respect to the weight w , i.e. the inner product $(\cdot | \cdot)_H$ on H is given by

$$(u|v)_H = \int_0^1 u(\rho) \overline{v(\rho)} w(\rho) d\rho$$

and as usual we write $\|\cdot\|_H := \sqrt{(\cdot | \cdot)_H}$. We set $\mathcal{D}(A_n) := \{u \in H : u, pu' \in AC_{\text{loc}}(0, 1), a_n u \in H\}$ and $A_n u := a_n u$ for $u \in \mathcal{D}(A_n)$.

Lemma 4.1. *The operator $A_n : \mathcal{D}(A_n) \subset H \rightarrow H$ is self-adjoint.*

Proof. Consider solutions of $a_n u = 0$. Since the coefficients of this linear ordinary differential equation are meromorphic and $\rho = 0$ as well as $\rho = 1$ are regular singular points, we can apply Frobenius' method to obtain series representations for solutions around $\rho = 0$ and $\rho = 1$. Around $\rho = 0$ the *Frobenius indices* are -2 and 1 , i.e. only one nontrivial solution belongs to H . According to the Weyl alternative we conclude that $\rho = 0$ is in the limit-point case (cf. e.g. [20]). Similarly, around $\rho = 1$ the indices are both $\frac{1}{2}$ and thus, neither of the two linearly independent solutions belongs to H which shows that $\rho = 1$ is in the limit-point case as well. It follows that the maximal operator A_n associated to a_n is self-adjoint (cf. [20]). \square

4.2 Factorization of A_0

Very similar to the energy critical case [10] there exists a factorization of the linearized operator A_0 that immediately implies its nonnegativity. To see this, note first that the potential term in eq. (5) for the ground state f_0 can be rewritten according to

$$\cos(2f_0(\rho)) = \cos(4 \arctan \rho) = \frac{1 - 6\rho^2 + \rho^4}{(1 + \rho^2)^2}.$$

Now we define two first-order operators B and \hat{B} on H by $\mathcal{D}(B) := \mathcal{D}(\hat{B}) := C_c^\infty(0, 1)$ as well as

$$Bu(\rho) := (1 - \rho^2)u'(\rho) - \frac{1 - 3\rho^2}{\rho(1 + \rho^2)}u(\rho)$$

and

$$\hat{B}u(\rho) := -(1 - \rho^2)u'(\rho) - \frac{3 - \rho^2}{\rho(1 + \rho^2)}u(\rho).$$

A simple integration by parts shows that $(Bu|v)_H = (u|\hat{B}v)_H$ for all $u, v \in C_c^\infty(0, 1)$ and thus, B and \hat{B} are adjoint to each other. Furthermore, by a direct computation one verifies that $A_0 u = \hat{B}B u$ for all $u \in C_c^\infty(0, 1)$ which is the desired factorization. We immediately obtain $(A_0 u|u)_H = (\hat{B}B u|u)_H = (B u|B u)_H \geq 0$ for all $u \in C_c^\infty(0, 1)$. Since A_0 is limit-point at both endpoints it follows that $C_c^\infty(0, 1)$ is a core for A_0 and the estimate $(A_0 u|u) \geq 0$ extends to all $u \in \mathcal{D}(A_0)$ by a density argument. Thus, A_0 is nonnegative and we have $\sigma(A_0) \subset [0, \infty)$.

4.3 The Spectrum of A_n

Proposition 4.1. *The point spectrum $\sigma_p(A_n)$ of A_n consists of exactly n negative eigenvalues, i.e. $\sigma_p(A_n) = \{\lambda_1, \dots, \lambda_n : 0 > \lambda_1 > \dots > \lambda_n\}$. Furthermore, the essential spectrum $\sigma_e(A_n)$ is given by $\sigma_e(A_n) = [0, \infty)$.*

Proof. Consider the equation $(\lambda - a_n)u = 0$. The Frobenius indices around $\rho = 1$ are $\frac{1 \pm \sqrt{-\lambda}}{2}$. Thus, for $\lambda \geq 0$ neither of the two linearly independent solutions belongs to H which shows $\sigma_p(A_n) \subset (-\infty, 0)$. Furthermore, we conclude $\sigma_e(A_n) = [0, \infty)$ by [18], p. 165, Theorem 11.5c.

As already discussed, the function θ_n defined by $\theta_n(\rho) := \rho\sqrt{1 - \rho^2}f_n'(\rho)$ is a solution of $a_n u = 0$. Bizoń [2] has shown that $f_n(\rho) \in (0, \pi)$ for $\rho \in (0, 1]$ and thus, if $f_n > \frac{\pi}{2}$, it can only have one maximum since

$$f_n''(\rho) = \frac{\sin(2f_n(\rho))}{\rho^2(1 - \rho^2)} < 0$$

for $f_n'(\rho) = 0$ (cf. eq. (3)). Similarly, if $f_n < \frac{\pi}{2}$, it can only have one minimum. Hence, the number of extrema of f_n equals the number of intersections of f_n with the line $\frac{\pi}{2}$ on $(0, 1)$ and therefore, θ_n has exactly n zeros on $(0, 1)$.

Applying the transformation $v(x) := \sinh(x)u(\tanh x)$, the equation $(\lambda - a_n)u = 0$ reduces to the eigenvalue problem

$$\lambda v(x) + v''(x) - \frac{2}{x^2}v(x) - Q_n(x)v(x) = 0 \tag{7}$$

where

$$Q_n(x) := \frac{2 \cos(2f_n(\tanh x))}{\sinh^2 x} - \frac{2}{x^2}.$$

Consequently, the function $\varphi_n(x) := \sinh(x)\theta_n(\tanh x)$ is a solution of eq. (7) with $\lambda = 0$ which has exactly n zeros on $x \in (0, \infty)$. Observe that the mapping $u \mapsto v$ is an isometry from $L_w^2(0, 1)$ to $L^2(0, \infty)$ and thus, the study of A_n is equivalent to the spectral problem for the $\ell = 1$ radial Schrödinger operator on $L^2(0, \infty)$ with potential Q_n . Note that $Q_n \in L^\infty(0, \infty)$ and from the Frobenius analysis of a_n it follows that the asymptotic behaviour of nontrivial solutions of eq. (7) is the same as if $Q_n \in C_c^\infty(0, \infty)$. Thus, the results of [9], XIII.3.B apply and we conclude that A_n has exactly n eigenvalues below the essential spectrum. \square

Remark 4.1. This observation has already been made by Bizoń [4]. Note further that θ_n is not an eigenfunction since it does not belong to H !

Remark 4.2. Negative eigenvalues of A_n lead to unstable mode solutions of eq. (5), i.e. $\phi(\sigma, \rho) := e^{(1+\sqrt{-\lambda})\sigma}v(\rho)$ is a solution of eq. (5) if $\lambda < 0$ belongs to $\sigma_p(A_n)$ and v is the associated eigenfunction. Thus, the n -th self-similar solution has exactly n unstable modes.

5 Well-Posedness of the Linearized Problem and Growth of Solutions

We consider the linear Cauchy problem

$$\begin{cases} \ddot{u}(\sigma) + A_0 u(\sigma) = 0 & \text{for } \sigma > 0 \\ u(0) = u_0, \dot{u}(0) = u_1 \end{cases} \quad (8)$$

for a function $u : [0, \infty) \rightarrow H$ and $u_0, u_1 \in \mathcal{D}(A_0)$ where $\dot{\cdot} := \frac{d}{d\sigma}$ and differentiation with respect to σ is understood in the strong sense. Eq. (8) is an operator formulation of eq. (6) with $f_n = f_0$ and thus, it describes the (rescaled) linear flow around the self-similar wave map f_0 .

According to the spectral theorem for self-adjoint operators (see e.g. [19]) there exists a spectral family $\{E(\lambda) : \lambda \in \mathbb{R}\}$ of orthogonal projections associated to A_0 such that for $u \in \mathcal{D}(A_0)$ and $v \in H$ we have

$$(A_0 u|v)_H = \int \text{id}_{\mathbb{R}} d\mu_{u,v}$$

where $\mu_{u,v} := \frac{1}{4}(\mu_{u+v} - \mu_{u-v} + i\mu_{u-iv} - i\mu_{u+iv})$ and μ_u is the unique measure defined on the Borel- σ -algebra on \mathbb{R} that satisfies $\mu_u((\lambda_1, \lambda_2]) = \|(E(\lambda_2) - E(\lambda_1))u\|_H^2$ for $u \in H$. Moreover, with the help of the spectral family $\{E(\lambda)\}$ it is possible to define functions of self-adjoint operators. Let $f : \mathbb{R} \rightarrow \mathbb{C}$ be measurable and set $\mathcal{D}(f(A_0)) := \{u \in H : \int |f|^2 d\mu_u < \infty\}$, $(f(A_0)u|v)_H := \int f d\mu_{u,v}$ for any $u \in \mathcal{D}(f(A_0))$ and $v \in H$. Note that in fact only the values of f on the spectrum of A_0 contribute. This yields a well-defined linear operator $f(A_0) : \mathcal{D}(f(A_0)) \subset H \rightarrow H$ which behaves nicely (see [19] for details and a functional calculus).

Proposition 5.1. *Let $u_0, u_1 \in \mathcal{D}(A_0)$. Then, the Cauchy problem eq. (8) has a unique solution $u : [0, \infty) \rightarrow \mathcal{D}(A_0) \subset H$ and the real-valued function $\sigma \mapsto \|A_0^{1/2}u(\sigma)\|_H^2 + \|\dot{u}(\sigma)\|_H^2$ is constant for all $\sigma \geq 0$.*

Proof. We set

$$g_\sigma(\lambda) := \begin{cases} \sin(\sigma\lambda)\lambda^{-1} & \text{for } \lambda > 0 \\ \sigma & \text{for } \lambda \leq 0 \end{cases}$$

and define $u(\sigma) := \cos(\sigma A_0^{1/2})u_0 + g_\sigma(A_0^{1/2})u_1$ via the functional calculus. Note that $A_0^{1/2}$ is well-defined since $\sigma(A_0) = [0, \infty)$. Applying Lebesgue's theorem on dominated convergence one immediately concludes that $\ddot{u}(\sigma) + A_0 u(\sigma) = 0$ and clearly, u satisfies the initial conditions. Furthermore, it follows easily that $u(\sigma)$ and $\dot{u}(\sigma)$ stay in $\mathcal{D}(A_0)$ for all $\sigma > 0$. A direct computation (functional calculus) yields $\|A_0^{1/2}u(\sigma)\|_H^2 + \|\dot{u}(\sigma)\|_H^2 = \|A_0^{1/2}u_0\|_H^2 + \|u_1\|_H^2$ which implies uniqueness as well. \square

Note that $(u_0, u_1) \mapsto \|A_0^{1/2}u_0\|_H^2 + \|u_1\|_H^2$ defines a norm on $\mathcal{D}(A_0^{1/2}) \times H$ since $0 \notin \sigma_p(A_0^{1/2})$ and hence, there do not exist growing solutions of eq. (8) which is the desired result. Thus, in view of the transformation $\phi \mapsto \tilde{\phi}$ we have shown that there do not exist linear perturbations of f_0 that grow faster than the gauge instability.

6 Discussion

We give a heuristic explanation why one cannot go any further in this formulation. This is most easily seen within a first-order formulation. Let $u : [0, \infty) \rightarrow H$ be the unique solution of eq. (8) with initial data $u_0, u_1 \in \mathcal{D}(A_0)$ and set $\phi(\sigma, \cdot) := e^\sigma u(\sigma)$. Then, ϕ satisfies the first-order equation

$$\frac{d}{d\sigma} \mathbf{u}(\sigma) = L \mathbf{u}(\sigma)$$

where $\mathbf{u}(\sigma) := (\phi(\sigma, \cdot), \phi_\sigma(\sigma, \cdot))$ and

$$L := \begin{pmatrix} 1 & 1 \\ -A_0 & 1 \end{pmatrix}.$$

ϕ (resp. \mathbf{u}) describes the linear flow around the self-similar wave map f_0 . Consider the operator L on the Banach space $X := \mathcal{D}(A_0^{1/2}) \times H$ with norm $\|(u_1, u_2)\|_X^2 := \|u_1\|_{A_0^{1/2}}^2 + \|u_2\|_H^2$ and $\mathcal{D}(L) := \mathcal{D}(A_0) \times \mathcal{D}(A_0^{1/2})$ where $\|\cdot\|_{A_0^{1/2}}$ denotes the graph norm of $A_0^{1/2}$. Then, it is easily seen that the spectrum of L can be calculated from the spectrum of A_0 and it is given by $\sigma(L) = \{z \in \mathbb{C} : -(z-1)^2 \in \sigma(A_0)\}$ which shows that $\sigma(L) = \{z \in \mathbb{C} : \operatorname{Re}(z) = 1\}$. We have already discussed the gauge instability which corresponds to the point $1 \in \sigma(L)$. Thus, the origin of the point $1 \in \sigma(L)$ is well-understood and in fact, we are only interested in stability of f_0 "modulo this gauge instability". If $1 \in \sigma(L)$ would be an isolated point one could define a projection operator to remove this instability. However, the rest of the spectrum makes this impossible and in what follows we give an intuitive explanation where it emanates from. The reason for this is a serious defect of the coordinate system (σ, ρ) . As already mentioned, it covers only the interior of the backward lightcone of the point $(T, 0)$. Thus, an outgoing wave packet can never leave the backward lightcone but cumulates near $\rho = 1$ as σ increases. This corresponds to an apparent instability which shows up in the spectrum of L . Unfortunately, one cannot go around this difficulty unless one changes coordinates which destroys the self-adjoint character of the problem. Thus, the result that there do not exist linear perturbations of f_0 that grow faster than the gauge instability is the best one can have in this formulation but still, it strongly supports the conjecture of stability of the wave map f_0 .

7 The Operators A_n for $n \rightarrow \infty$

The self-similar wave maps f_n can be constructed numerically using a shooting and matching technique [1]. By the same method one may calculate the point spectrum of A_n (cf. [4]). Results for A_1, \dots, A_4 are given in Table 1 where $\mu_j := \sqrt{-\lambda_j}$.

7.1 The Operator A_∞

We emphasize the remarkable fact that the horizontal rows in Table 1 seem to converge. The question is if one can construct an operator A_∞ such that the point spectrum of A_∞ is the limit of the point spectra of A_n for $n \rightarrow \infty$. Since $f_n \rightarrow f_\infty \equiv \frac{\pi}{2}$ pointwise on $(0, 1]$, the obvious strategy is to investigate linear perturbations around f_∞ described by eq. (6) with $f_n = f_\infty$. Thus, we consider the Sturm-Liouville expression

$$a_\infty u := \frac{1}{w} (-(pu)') + q_\infty u$$

where $q_\infty(\rho) := \frac{-2+\rho^2}{(1-\rho^2)^2}$. Again, we use Frobenius' method to classify the endpoints. The behaviour of solutions of $a_\infty u = 0$ around $\rho = 1$ is the same as before for a_n and thus, $\rho = 1$ is in the limit-point case again. However, around $\rho = 0$ the Frobenius indices are $\frac{-1 \pm i\sqrt{7}}{2}$ and thus, all solutions of $a_\infty u = 0$ belong to H near $\rho = 0$. According to the Weyl alternative the point $\rho = 0$ is in the limit-circle case and hence, we have to specify a boundary condition of the form $[u, \chi]_\rho(0) = 0$ for a fixed function χ in order to obtain a self-adjoint operator (cf. [20]). The question is how to choose χ . Bizoň [3] has studied an analogous problem for a Yang-Mills system. Following his proposal we choose χ in such a way that $a_\infty u = 0$ has a nontrivial solution just like the gauge instability for a_n . Thus, we denote by $\tilde{\chi}$ a real-valued nontrivial function satisfying $a_\infty \tilde{\chi} = 0$ with asymptotic behaviour $\tilde{\chi}(\rho) \sim (1-\rho)^{1/2}$ for $\rho \rightarrow 1$. Such a function exists by Frobenius' method and it is unique up to constant multiples since the

second linearly independent solution around $\rho = 1$ contains a log term. Using an appropriate smooth cut-off function we construct a χ that coincides with $\tilde{\chi}$ on $(0, \frac{1}{2})$ and satisfies $\chi, a_\infty \chi \in H$ as well as $\chi, p\chi' \in AC_{\text{loc}}(0, 1)$. Then, we define

$$\mathcal{D}(A_\infty) := \{u \in H : u, pu' \in AC_{\text{loc}}(0, 1), a_\infty u \in H, [u, \chi]_p(0) = 0\}$$

and $A_\infty u := a_\infty u$ for $u \in \mathcal{D}(A_\infty)$. Invoking Sturm–Liouville theory [20] we conclude that $A_\infty : \mathcal{D}(A_\infty) \subset H \rightarrow H$ is self-adjoint.

7.2 The Point Spectrum of A_∞

Since the asymptotic behaviour of solutions of $(\lambda - a_\infty)u = 0$ around $\rho = 1$ is the same as for $(\lambda - a_n)u = 0$, we conclude that $\sigma_p(A_\infty) \subset (-\infty, 0)$. Thus, we restrict ourselves to $\lambda < 0$. The eigenvalue equation $(\lambda - a_\infty)u = 0$ is a linear ordinary differential equation of second order with four regular singular points $\rho = 0, \pm 1, \infty$. However, applying the transformation $\rho \mapsto z := \rho^2$ we can reduce this number by 1 and we are left with the three regular singular points $z = 0, 1, \infty$. Thus, solutions are given in terms of hypergeometric functions (cf. [5]). We define a new unknown

$$v(z) := z^\alpha (1 - z)^\beta u(\sqrt{z})$$

where $\alpha := -\frac{1}{4}(-1 + i\sqrt{7})$ and $\beta := -\frac{1}{2}(1 + \sqrt{-\lambda})$. The equation $(\lambda - a_\infty)u = 0$ transforms into

$$z(1 - z)v'' + [c - (a + b + 1)z]v' - abv = 0$$

where $a := \frac{1}{4}(1 + 2\sqrt{-\lambda} + i\sqrt{7})$, $b := \frac{1}{4}(3 + 2\sqrt{-\lambda} + i\sqrt{7})$ and $c := 1 + \frac{i\sqrt{7}}{2}$. The solution $v_1(\cdot, \lambda)$ defined by $v_1(z, \lambda) := {}_2F_1(a, b; a + b + 1 - c; 1 - z)$ corresponds to the solution of $(\lambda - a_\infty)u = 0$ which belongs to H near $\rho = 1$. v_1 can be written as a linear combination of the form

$$v_1(z, \lambda) \propto {}_2F_1(a, b; c; z) + m(\lambda)z^{1-c}{}_2F_1(a + 1 - c, b + 1 - c; 2 - c; z)$$

where the connection coefficient $m(\lambda)$ can be given in terms of the Γ -function [5] and reads

$$m(\lambda) = \frac{\Gamma(a + 1 - c)\Gamma(b + 1 - c)\Gamma(c - 1)}{\Gamma(a)\Gamma(b)\Gamma(1 - c)}.$$

Thus, the boundary condition at $\rho = 0$ translates into $m(\lambda) = m(0)$. This transcendental equation can be solved numerically and the results are given in Table 1.

μ_n	A_1	A_2	A_3	A_4	A_∞
μ_1	5.333625	5.304	5.30	5.3	5.3009
μ_2		58.0701	57.68	57.6	57.637
μ_3			625	620	619.61

Table 1: Approximate eigenvalues

Thus, as conjectured, the point spectrum of A_∞ seems to be the limit of the point spectra of A_n .

8 Generalization to higher equivariance indices

Finally, we consider more general equivariant wave maps described by the equation

$$\psi_{tt} - \psi_{rr} - \frac{2}{r}\psi_r + \frac{\ell(\ell + 1)}{2} \frac{\sin(2\psi)}{r^2} = 0 \quad (9)$$

where $\ell \in \mathbb{N}$ is the equivariance index (see e.g. [12] for a derivation of this equation). Self-similar solutions $\psi(t, r) = f(\frac{r}{T-t})$ of eq. (9) satisfy

$$f'' + \frac{2}{\rho}f' - \frac{\ell(\ell + 1)}{2} \frac{\sin(2f)}{\rho^2(1 - \rho^2)} = 0$$

and again, we require $f(0) = 0$ and $f(1) = \frac{\pi}{2}$ as regularity conditions. Bizoń's proof of the existence of a countable set $\{f_{n,\ell} : n = 0, 1, 2, \dots\}$ of self-similar solutions carries over to this more general situation (see [2]). In fact, the solutions $f_{n,\ell}$ have analogous properties for all ℓ . The index n counts the number of intersections of $f_{n,\ell}$ with the line $\frac{\pi}{2}$ in the interval $\rho \in [0, 1)$ and we have $f_{n,\ell}(\rho) \in (0, \pi)$ for all n and $\rho \in (0, 1]$. Furthermore, the asymptotic behaviour of $f_{n,\ell}$ for $\rho \rightarrow 0$ is $f_{n,\ell}(\rho) \sim \rho^\ell$.

Linear perturbations ϕ around $f_{n,\ell}$ satisfy the evolution equation

$$\phi_{\sigma\sigma} - 2\phi_\sigma - (1 - \rho^2)^2 \phi_{\rho\rho} - \frac{2(1 - \rho^2)^2}{\rho} \phi_\rho + \frac{\ell(\ell + 1)(1 - \rho^2) \cos(2f_{n,\ell})}{\rho^2} \phi = 0$$

and analogous to the case $\ell = 1$ we define the Sturm-Liouville expressions $a_{n,\ell}$ by

$$a_{n,\ell} u := \frac{1}{w} (-(pu)') + q_{n,\ell} u$$

where $w(\rho) := \frac{\rho^2}{(1 - \rho^2)^2}$, $p(\rho) := \rho^2$ and

$$q_{n,\ell}(\rho) := \frac{\ell(\ell + 1)(1 - \rho^2) \cos(2f_{n,\ell}(\rho)) - \rho^2}{(1 - \rho^2)^2}.$$

The Frobenius indices for the equation $a_{n,\ell} u = 0$ are $\{-\ell - 1, \ell\}$ at $\rho = 0$ and $\{\frac{1}{2}, \frac{1}{2}\}$ at $\rho = 1$ and thus, both endpoints are in the limit-point case. It follows that the associated maximal operator $A_{n,\ell}$ on $H = L^2_w(0, 1)$ is self-adjoint. Again, the study of $A_{n,\ell}$ is equivalent to the spectral problem for the radial Schrödinger operator and thus, the rest of the analysis remains unchanged. Hence, we have the following theorem.

Theorem 8.1. *The spectrum of $A_{n,\ell}$ consists of a continuous part $\sigma_e(A_{n,\ell}) = [0, \infty)$ and exactly n negative eigenvalues. As a consequence, $A_{0,\ell}$ is nonnegative, the Cauchy problem*

$$\begin{cases} \ddot{u}(\sigma) + A_{0,\ell} u(\sigma) = 0 \text{ for } \sigma > 0 \\ u(0) = u_0, \dot{u}(0) = u_1 \end{cases}$$

for $u_0, u_1 \in \mathcal{D}(A_{0,\ell})$ is well-posed and the real-valued function $\sigma \mapsto \|A_{0,\ell}^{1/2} u(\sigma)\|_H^2 + \|\dot{u}(\sigma)\|_H^2$ is constant for all $\sigma \geq 0$.

9 Acknowledgments

The authors would like to thank Piotr Bizoń and Horst R. Beyer for helpful discussions. This work has been supported by the Austrian Fonds zur Förderung der wissenschaftlichen Forschung FWF Project P19126 and partly by the Fundacion Federico.

References

- [1] Stefan Åminneborg and Lars Bergström. On self-similar global textures in an expanding universe. *Phys. Lett. B*, 362(1-4):39–45, 1995.
- [2] Piotr Bizoń. Equivariant self-similar wave maps from Minkowski spacetime into 3-sphere. *Comm. Math. Phys.*, 215(1):45–56, 2000.
- [3] Piotr Bizoń. Formation of singularities in Yang-Mills equations. *Acta Phys. Polon. B*, 33(7):1893–1922, 2002.
- [4] Piotr Bizoń, Tadeusz Chmaj, and Zbislav Tabor. Dispersion and collapse of wave maps. *Nonlinearity*, 13(4):1411–1423, 2000.
- [5] Arthur Erdélyi, Wilhelm Magnus, Fritz Oberhettinger, and Francesco G. Tricomi. *Higher transcendental functions. Vols. I, II*. McGraw-Hill Book Company, Inc., New York-Toronto-London, 1953. Based, in part, on notes left by Harry Bateman.
- [6] Sergiu Klainerman and Sigmund Selberg. Bilinear estimates and applications to nonlinear wave equations. *Commun. Contemp. Math.*, 4(2):223–295, 2002.

- [7] J. Krieger, W. Schlag, and D. Tataru. Renormalization and blow up for charge one equivariant critical wave maps. *Invent. Math.*, 171(3):543–615, 2008.
- [8] Frank Merle and Hatem Zaag. Determination of the blow-up rate for the semilinear wave equation. *Amer. J. Math.*, 125(5):1147–1164, 2003.
- [9] Michael Reed and Barry Simon. *Methods of modern mathematical physics. IV. Analysis of operators*. Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1978.
- [10] Igor Rodnianski and Jacob Sterbenz. On the formation of singularities in the critical $o(3)$ sigma-model, 2006. Preprint arXiv.org:math/0605023.
- [11] Jalal Shatah. Weak solutions and development of singularities of the $SU(2)$ σ -model. *Comm. Pure Appl. Math.*, 41(4):459–469, 1988.
- [12] Jalal Shatah and A. Shadi Tahvildar-Zadeh. On the Cauchy problem for equivariant wave maps. *Comm. Pure Appl. Math.*, 47(5):719–754, 1994.
- [13] Thomas C. Sideris. Global existence of harmonic maps in Minkowski space. *Comm. Pure Appl. Math.*, 42(1):1–13, 1989.
- [14] Michael Struwe. Equivariant wave maps in two space dimensions. *Comm. Pure Appl. Math.*, 56(7):815–823, 2003. Dedicated to the memory of Jürgen K. Moser.
- [15] Terence Tao. Global regularity of wave maps. II. Small energy in two dimensions. *Comm. Math. Phys.*, 224(2):443–544, 2001.
- [16] Daniel Tataru. The wave maps equation. *Bull. Amer. Math. Soc. (N.S.)*, 41(2):185–204 (electronic), 2004.
- [17] N. Turok and D. Spergel. Global texture and the microwave background. *Phys. Rev. Lett.*, 64:2736–2739, 1990.
- [18] Joachim Weidmann. *Spectral theory of ordinary differential operators*, volume 1258 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1987.
- [19] Kōsaku Yosida. *Functional analysis*. Classics in Mathematics. Springer-Verlag, Berlin, 1995. Reprint of the sixth (1980) edition.
- [20] Anton Zettl. *Sturm-Liouville theory*, volume 121 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2005.